

THE SMALLEST HYPERBOLIC 6-MANIFOLDS

BRENT EVERITT, JOHN RATCLIFFE, AND STEVEN TSCHANTZ

ABSTRACT. By gluing together copies of an all-right angled Coxeter polytope a number of open hyperbolic 6-manifolds with Euler characteristic -1 are constructed. They are the first known examples of hyperbolic 6-manifolds having the smallest possible volume.

1. INTRODUCTION

The last few decades has seen a surge of activity in the study of finite volume hyperbolic manifolds—that is, complete Riemannian n -manifolds of constant sectional curvature -1 . Not surprisingly for geometrical objects, volume has been, and continues to be, the most important invariant for understanding their sociology. The possible volumes in a fixed dimension forms a well-ordered subset of $\mathbb{R}$, indeed a discrete subset except in 3-dimensions (where the orientable manifolds at least have ordinal type ω^ω). Thus it is a natural problem with a long history to construct examples of manifolds with minimal volume in a given dimension.

In 2-dimensions the solution is classical, with the minimum volume in the compact orientable case achieved by a genus 2 surface, and in the non-compact orientable case by a once-punctured torus or thrice-punctured sphere (the identities of the manifolds are of course also known in the non-orientable case). In 3-dimensions the compact orientable case remains an open problem with the Matveev-Fomenko-Weeks manifold [16, 30] obtained via $(5, -2)$ -Dehn surgery on the the sister of the figure-eight knot complement conjecturally the smallest. Amongst the non-compact orientable 3-manifolds the figure-eight knot complement realizes the minimum volume [17] and the Gieseking manifold (obtained by identifying the sides of a regular hyperbolic tetrahedron as in [14, 20]) does so for the non-orientable ones [1]. One could also add “arithmetic” to our list of adjectives and so have eight optimization problems to play with (so that the Matveev-Fomenko-Weeks manifold is known to be the minimum volume orientable, arithmetic compact 3-manifold, see [5]).

When $n \geq 4$ the picture is murkier, although in even dimensions we have recourse to the Gauss-Bonnet Theorem, so that in particular the minimal volume a $2m$ -dimensional hyperbolic manifold could possibly have is when the Euler characteristic χ satisfies $|\chi| = 1$. The first examples of non-compact 4-manifolds with $\chi = 1$ were constructed in [23] (see also [13]). The compact case remains a difficult unsolved problem, although if we restrict to arithmetic manifolds, then it is known [2,8] that a minimal volume arithmetic compact orientable 4-manifold M

1991 *Mathematics Subject Classification.* Primary 57M50.

The first author is grateful to the Mathematics Department, Vanderbilt University for its hospitality during a stay when the results of this paper were obtained.

has $\chi \leq 16$ and M is isometric to the orbit space of a torsion-free subgroup of the hyperbolic Coxeter group $[5, 3, 3, 3]$. The smallest compact hyperbolic 4-manifold currently known to exist has $\chi = 8$ and is constructed in [8]. Manifolds of very small volume have been constructed in 5-dimensions [13, 24], but the smallest volume 6-dimensional example hitherto known has $\chi = -16$ [13].

In this paper we announce the discovery of a number of non-compact non-orientable hyperbolic 6-manifolds with Euler characteristic $\chi = -1$. The method of construction is classical in that the manifolds are obtained by identifying the sides of a 6-dimensional hyperbolic Coxeter polytope.

2. COXETER POLYTOPES


Let C be a convex (not necessarily bounded) polytope of finite volume in a simply connected space X^n of constant curvature. Call C a Coxeter polytope if the dihedral angle subtended by two intersecting $(n-1)$ -dimensional sides is π/m for some integer $m \geq 2$. When $X^n = S^n$ or the Euclidean space E^n , such polyhedra have been completely classified [11], but in the hyperbolic space H^n , a complete classification remains a difficult problem (see for example [27] and the references there).

If Γ is the group generated by reflections in the $(n-1)$ -dimensional sides of C , then Γ is a discrete cofinite subgroup of the Lie group $\text{Isom } X^n$, and every discrete cofinite reflection group in $\text{Isom } X^n$ arises in this way from some Coxeter polytope, which is uniquely defined up to isometry. The Coxeter symbol for C (or Γ) has nodes indexed by the $(n-1)$ -dimensional sides, and an edge labeled m joining the nodes corresponding to sides that intersect with angle π/m (label the edge joining the nodes of non-intersecting sides by ∞). In practice the labels 2 and 3 occur often, so that edges so labeled are respectively removed or left unlabeled.

Let Λ be a $(n+1)$ -dimensional Lorentzian lattice, that is, an $(n+1)$ -dimensional free $\mathbb{Z}$ -module equipped with a $\mathbb{Z}$ -valued bilinear form of signature $(n, 1)$. For each n , there is a unique such Λ , denoted $I_{n,1}$, that is odd and self-dual (see [25, Theorem V.6], or [18, 19]). By [4], the group $O_{n,1}\mathbb{Z}$ of automorphisms of $I_{n,1}$ acts discretely, cofinitely by isometries on the hyperbolic space H^n obtained by projectivising the negative norm vectors in the Minkowski space-time $I_{n,1} \otimes \mathbb{R}$ (to get a faithful action one normally passes to the centerless version $PO_{n,1}\mathbb{Z}$).

Vinberg and Kaplinskaja showed [28, 29] that the subgroup Reflec_n of $PO_{n,1}\mathbb{Z}$ generated by reflections in positive norm vectors has finite index if and only if $n \leq 19$, thus yielding a family of cofinite reflection groups and corresponding finite volume Coxeter polytopes in the hyperbolic spaces H^n for $2 \leq n \leq 19$. Indeed, Conway and Sloane have shown [9, Chapter 28] or [10], that for $n \leq 19$ the quotient of $PO_{n,1}\mathbb{Z}$ by Reflec_n is a subgroup of the automorphism group of the Leech lattice. Borcherds [3] showed that the (non self-dual) even sublattice of $I_{21,1}$ also acts cofinitely, yielding the highest dimensional example known of a Coxeter group acting cofinitely on hyperbolic space.

When $4 \leq n \leq 9$ the group $\Gamma = \text{Reflec}_n$ has Coxeter symbol,



with $n + 1$ nodes and C a non-compact, finite volume n -simplex Δ^n (when $n > 9$, the polytope C has a more complicated structure).

Let v be the vertex of Δ^n opposite the side F_1 marked on the symbol, and let Γ_v be the stabilizer in Γ of this vertex. This stabilizer is also a reflection group with symbol as shown, and is finite for $4 \leq n \leq 8$ (being the Weyl group of type A_4, D_5, E_6, E_7 and E_8 respectively) and infinite for $n = 9$ (when it is the affine Weyl group of type $\tilde{E}_8$). Let

$$P_n = \bigcup_{\gamma \in \Gamma_v} \gamma(\Delta^n),$$

a convex polytope obtained by gluing $|\Gamma_v|$ copies of the simplex Δ^n together. Thus, P_n has finite volume precisely when $4 \leq n \leq 8$, although it is non-compact, with a mixture of finite vertices in H^n and cusped ones on ∂H^n . In any case, P_n is an all right-angled Coxeter polytope: its sides meet with dihedral angle $\pi/2$ or are disjoint. This follows immediately from the observation that the sides of P_n arise from the Γ_v -images of the side of Δ^n opposite v , and this side intersects the other sides of Δ^n in dihedral angles $\pi/2$ or $\pi/4$. Vinberg has conjectured that $n = 8$ is the highest dimension in which finite volume all-right angled polytopes exist in hyperbolic space.

The volume of the polytope P_n is given by

$$\text{vol}(P_n) = |\Gamma_v| \text{vol}(\Delta^n) = |\Gamma_v| [\text{PO}_{n,1}\mathbb{Z} : \Gamma] \text{covol}(\text{PO}_{n,1}\mathbb{Z}),$$

where $\text{covol}(\text{PO}_{n,1}\mathbb{Z})$ is the volume of a fundamental region for the action of $\text{PO}_{n,1}\mathbb{Z}$ on H^n (and for $4 \leq n \leq 9$ the index $[\text{PO}_{n,1}\mathbb{Z} : \Gamma] = 1$). When n is even, we have by [26] and [22],

$$\text{covol}(\text{PO}_{n,1}\mathbb{Z}) = \frac{(2^{\frac{n}{2}} \pm 1) \pi^{\frac{n}{2}}}{n!} \prod_{k=1}^{\frac{n}{2}} |B_{2k}|,$$

with B_{2k} the $2k$ -th Bernoulli number and with the plus sign if $n \equiv 0, 2 \pmod{8}$ and the minus sign otherwise.

Alternatively (when n is even), we have recourse to the Gauss-Bonnet Theorem, so that $\text{vol}(P_n) = \kappa_n |\Gamma_v| \chi(\Gamma)$, where $\chi(\Gamma)$ is the Euler characteristic of the Coxeter group Γ and $\kappa_n = 2^n (n!)^{-1} (-\pi)^{n/2} (n/2)!$. The Euler characteristic of Coxeter groups can be easily computed from their symbol (see [6, 7] or [13, Theorem 9]). Indeed, when $n = 6$, $\chi(\Gamma) = -1/\mathcal{L}$ where $\mathcal{L} = 2^{10} 3^4 5$ and so $\text{vol}(P_6) = 8\pi^3 |E_6| / 15\mathcal{L} = \pi^3 / 15$.

The Coxeter symbol for P_n has a nice description in terms of finite reflection groups. If v' is the vertex of Δ^n opposite the side F_2 , let Γ_e be the pointwise stabilizer of $\{v, v'\}$: the elements thus stabilize the edge e of Δ^n joining v and v' .

Now consider the Cayley graph $\mathcal{C}_v$ for Γ_v with respect to the generating reflections in the sides of the symbol for Γ_v . Thus, $\mathcal{C}_v$ has vertices in one to one correspondence with the elements of Γ_v and for each generating reflection s_α , an undirected edge labeled s_α connecting vertices γ_1 and γ_2 if and only if $\gamma_2 = \gamma_1 s_\alpha$ in Γ_v . In particular, $\mathcal{C}_v$ has s_2 labeled edges corresponding to the reflection in F_2 . Removing these s_2 -edges decomposes $\mathcal{C}_v$ into components, each of which is a copy of the Cayley graph $\mathcal{C}_e$ for Γ_e , with respect to the generating reflections.

Take as the nodes of the symbol for P_n these connected components. If two components have an s_2 -labeled edge running between any two of their vertices in $\mathcal{C}_v$, then leave the corresponding nodes unconnected, otherwise, connect them by

an edge labeled ∞ . The resulting symbol (respectively the polytope P_n) thus has $|\Gamma_v|/|\Gamma_e|$ nodes (resp. sides). The number of sides of P_n for $n = 4, 5, 6, 7, 8$ is 10, 16, 27, 56 and 240 respectively.

3. CONSTRUCTING THE MANIFOLDS

We now restrict our attention to the case $n = 6$. We work in the hyperboloid model of hyperbolic 6-space

$$H^6 = \{x \in \mathbb{R}^7 : x_1^2 + x_2^2 + \cdots + x_6^2 - x_7^2 = -1 \text{ and } x_7 > 0\}$$

and represent the isometries of H^6 by Lorentzian 7×7 matrices that preserve H^6 . The right-angled polytope P_6 has 27 sides each congruent to P_5 . We position P_6 in H^6 so that 6 of its sides are bounded by the 6 coordinate hyperplanes $x_i = 0$ for $i = 1, \dots, 6$ and these 6 sides intersect at the center e_7 of H^6 . Let K_6 be the group of 64 diagonal Lorentzian 7×7 matrices $\text{diag}(\pm 1, \dots, \pm 1, 1)$. The set $Q_6 = K_6 P_6$, which is the union of 64 copies of P_6 , is a right-angled convex polytope with 252 sides. We construct hyperbolic 6-manifolds, with $\chi = -8$, by gluing together the sides of Q_6 by a proper side-pairing with side-pairing maps of the form rk with k in K_6 and r a reflection in a side S of Q_6 . The side-pairing map rk pairs the side $S' = kS$ to S (see §11.1 and §11.2 of [21] for a discussion of proper side-pairings). We call such a side-pairing of Q_6 simple. We searched for simple side-pairings of Q_6 that yield a hyperbolic 6-manifold M with a freely acting $\mathbb{Z}/8$ symmetry group that permutes the 64 copies of P_6 making up M in such a way that the resulting quotient manifold is obtained by gluing together 8 copies of P_6 . Such a quotient manifold has $\chi = -8/8 = -1$. This is easier said than done, since the search space of all possible side-pairings of Q_6 is very large. We succeeded in finding desired side-pairings of Q_6 by employing a strategy that greatly reduces the search space. The strategy is to extend a side-pairing in dimension 5 with the desired properties to a side-pairing in dimension 6 with the desired properties.

Let $Q_5 = \{x \in Q_6 : x_1 = 0\}$. Then Q_5 is a right-angled convex 5-dimensional polytope with 72 sides. Note that Q_5 is the union $K_5 P_5$ of 32 copies of P_5 where $P_5 = \{x \in P_6 : x_1 = 0\}$ and K_5 is the group of 32 diagonal Lorentzian 7×7 matrices $\text{diag}(1, \pm 1, \dots, \pm 1, 1)$. A simple side-pairing of Q_6 that yields a hyperbolic 6-manifold M restricts to a simple side-pairing of Q_5 that yields a hyperbolic 5-manifold which is a totally geodesic hypersurface of M . All the orientable hyperbolic 5-manifolds that are obtained by gluing together the sides of Q_5 by a simple side-pairing are classified in [24].

We started with the hyperbolic 5-manifold N , numbered 27 in [24], obtained by gluing together the sides of Q_5 by the simple side-pairing with side-pairing code 2B7JB47JG81. The manifold N has a freely acting $\mathbb{Z}/8$ symmetry group that permutes the 32 copies of P_5 making up N in such a way that the resulting quotient manifold is obtained by gluing together 4 copies of P_5 . A generator of the $\mathbb{Z}/8$ symmetry group of N is represented by the Lorentzian 6×6 matrix

$$\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & -1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & -1 & 0 & 1 \\ -1 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}.$$

N	SP	H_1	H_2	H_3	H_4	H_5
		0248	0248	0248	0248	0248
1	GW8dNEEdN4ZJ01k211PIY	0401	1910	4821	1500	0000
2	HX9dNFEcM5aKU6f3f6UKa	0401	1810	8710	5500	0000
3	HX9dNFEcM5YIO113110IY	0401	2900	7810	4400	1000
4	HX9dNFEcM5YIO613160IY	0401	2800	7910	4400	1000
5	HX9dNFEcM5YIOx131y0IY	0211	2800	4821	1400	1000
6	HX9dNFEcM5YIOy131x0IY	0211	2800	4930	1400	1000
7	HX9dNFEcM5aKUxf3fyUKa	0301	1900	5630	2500	0000

TABLE 1. Side-pairing codes and homology groups of the seven examples

The strategy is to search for simple side-pairings of Q_6 that yield a hyperbolic 6-manifold with a freely acting $\mathbb{Z}/8$ symmetry group with generator represented by the following Lorentzian 7×7 matrix that extends the above Lorentzian 6×6 matrix.

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & -1 & 0 & 1 \\ 0 & -1 & 0 & -1 & -1 & 0 & 2 \end{pmatrix}$$

For such a side-pairing the resulting quotient manifold can be obtained by gluing together 8 copies of P_6 by a proper side-pairing. By a computer search we found 14 proper side-pairings of 8 copies of P_6 in this way, and hence we found 14 hyperbolic 6-manifolds with $\chi = -1$. Each of these 14 manifolds is noncompact with volume $8\text{vol}(P_6) = 8\pi^3/15$ and five cusps. These 14 hyperbolic 6-manifolds represent at least 7 different isometry types, since they represent 7 different homology types. Table 1 lists side-pairing codes for 7 simple side-pairings of Q_6 whose $\mathbb{Z}/8$ quotient manifold has homology groups isomorphic to $\mathbb{Z}^a \oplus (\mathbb{Z}/2)^b \oplus (\mathbb{Z}/4)^c \oplus (\mathbb{Z}/8)^d$ for nonnegative integers a, b, c, d encoded by $abcd$ in the table. In particular, all 7 manifolds in Table 1 have a finite first homology group.

All of our examples, with $\chi = -1$, can be realized as the orbit space H^6/Γ of a torsion-free subgroup Γ of $\text{PO}_{6,1}\mathbb{Z}$ of minimal index. These manifolds are the first examples of hyperbolic 6-manifolds having the smallest possible volume. All these manifolds are nonorientable. In the near future, we hope to construct orientable examples of noncompact hyperbolic 6-manifolds having $\chi = -1$.

REFERENCES

1. C Adams. *The non-compact hyperbolic 3-manifold of minimum volume*. Proc. AMS, 100 (1987), 601-606.
2. M Belolipetsky. *On volumes of arithmetic quotients of $SO(1, n)$* . Ann. Scuola Norm. Sup. Pisa Cl. Sci. (to appear).
3. R Borcherds. *Automorphism groups of Lorentzian lattices*. J. Algebra 111(1) (1987), 133-153.
4. A Borel and Harish-Chandra. *Arithmetic subgroups of algebraic groups*. Ann. of Math., 75(3) (1962), 485-535.
5. T Chinburg, E Friedman, K N Jones, and A W Reid. *The arithmetic hyperbolic 3-manifold of smallest volume*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 30 (2001), 1-40.

6. I M Chiswell. *The Euler characteristic of graph products and Coxeter groups*. in Discrete Groups and Geometry, W J Harvey and Colin Maclachlan (Editors), London Math. Soc. Lect. Notes, 173 (1992), 36-46.
7. I M Chiswell. *Euler characteristics of groups*. Math. Z., 147 (1976), 1-11.
8. M D E Conder and C Maclachlan. *Small volume compact hyperbolic 4-manifolds*. to appear, Proc. Amer. Math. Soc.
9. J Conway and N J A Sloane. *Sphere Packings, Lattices and Groups*. Second Edition, Springer 1993.
10. J Conway and N J A Sloane. *Leech roots and Vinberg groups*. Proc. Roy. Soc. London Ser. A 384 (no. 1787) (1982), 233-258.
11. H S M Coxeter. *Discrete groups generated by reflections*. Ann. of Math, 35(2) (1934), 588-621.
12. M W Davis. *A hyperbolic 4-manifold*. Proc. Amer. Math. Soc., 93 (1985), 325-328.
13. B Everitt. *Coxeter groups and hyperbolic manifolds*. Math. Ann. 330(1), (2004), 127-150.
14. B Everitt. *3-Manifolds from Platonic solids*. Top. App. 138 (2004), 253-263.
15. B Everitt, J Ratcliffe and S Tschantz. *Arithmetic hyperbolic 6-manifolds of smallest volume* (in preparation).
16. V Matveev and A Fomenko. *Constant energy surfaces of Hamilton systems, enumeration of three-dimensional manifolds in increasing order of complexity, and computation of volumes of closed hyperbolic manifolds*. Russian Math. Surveys, 43 (1988), 3-24.
17. C Cao and G R Meyerhoff. *The orientable cusped hyperbolic 3-manifolds of minimum volume*. Invent. Math. 146(3) (2001), 451-478.
18. J Milnor and D Husemoller. *Symmetric Bilinear Forms*. Springer, 1973.
19. A Neumaier and J J Seidel. *Discrete Hyperbolic Geometry*. Combinatorica 3 (1983), 219-237.
20. I Prok. *Classification of dodecahedral space forms*. Beiträge Algebra Geom. 39(2) (1998), 497-515.
21. J Ratcliffe. *Foundations of hyperbolic manifolds*. Graduate Texts in Mathematics 149, Springer 1994.
22. J Ratcliffe and S Tschantz. *Volumes of integral congruence hyperbolic manifolds*. J. Reine Angew. Math. 488 (1997), 55-78.
23. J Ratcliffe and S Tschantz. *The volume spectrum of hyperbolic 4-manifolds*. Experiment. Math. 9 (2000), no. 1, 101-125.
24. J Ratcliffe and S Tschantz. *Integral congruence two hyperbolic 5-manifolds*. Geom. Dedicata 107 (2004), 187-209.
25. J-P Serre. *A Course in Arithmetic*. Graduate Texts in Mathematics 7, Springer 1973.
26. C L Siegel. *Über die analytische Theorie der quadratischen Formen II*. Ann. Math. 37 (1936), 230-263.
27. E B Vinberg. *Hyperbolic Reflection Groups*. Uspekhi Mat. Nauk 40:1 (1985), 29-66, = Russian Math. Surveys 40:1 (1985), 31-75.
28. E B Vinberg and I M Kaplinskaja. *On the groups $O_{18,1}(\mathbb{Z})$ and $O_{19,1}(\mathbb{Z})$* . SMD 19(1) (1978), 194-197. = DAN 238(6) (1978), 1273-1275.
29. E B Vinberg. *On the groups of units of certain quadratic forms*. Sb. 16(1) (1972), 17-35, = Mat. Sb. 87(1) (1972), 18-36.
30. J Weeks. *Hyperbolic structures on 3-manifolds*. PhD thesis, Princeton University.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, ENGLAND
E-mail address: `bje1@york.ac.uk`

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240, USA
E-mail address: `ratclifj@math.vanderbilt.edu`

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240, USA
E-mail address: `tschantz@math.vanderbilt.edu`